



TEXTBOOK TESTBANKS

Solutions Manual for Precalculus with Limits 6th Larson

SOLUTIONS MANUAL SAMPLE PREVIEW

PUBLISHER

Cengage

COPYRIGHT

2027

ISBN

9798214407760

RESOURCE TYPE

Solutions Manual

*Solutions Manual for Precalculus with
Limits 6th Edition by Larson*

ISBN: 9798214407760

Solution and Answer Guide

LARSON, PRECALCULUS 2027, 9798214407050;

APPENDIX A: REVIEW OF FUNDAMENTAL CONCEPTS OF ALGEBRA

TABLE OF CONTENTS

Appendix A.1 Real Numbers and Their Properties	2146
Appendix A.2 Exponents and Radicals	2155
Appendix A.3 Polynomials and Factoring.....	2166
Appendix A.4 Rational Expressions	2179
Appendix A.5 Solving Equations	2193
Appendix A.6 Linear Inequalities in One Variable.....	2210
Appendix A.7 Errors and the Algebra of Calculus.....	2225

APPENDIX A.1 REAL NUMBERS AND THEIR PROPERTIES

A.1.1.

irrational

A.1.2.

Origin

A.1.3.

terms

A.1.4.

Zero-Factor Property

A.1.5.

Yes. $|3 - 10| = |-7| = 7$ and $|10 - 3| = |7| = 7$.

A.1.6.

- (a) iii
- (b) iv
- (c) ii
- (d) v
- (e) i

A.1.7.

$-9, -\frac{7}{2}, 5, \frac{2}{3}, \sqrt{3}, 0, 3, -4, 2, -11$

- (a) Natural numbers: 5, 8, 2
- (b) Whole numbers: 5, 0, 8, 2
- (c) Integers: $-9, 5, 0, 8, -4, 2, -11$
- (d) Rational numbers: $-9, -\frac{7}{2}, 5, \frac{2}{3}, 0, 8, -4, 2, -11$
- (e) Irrational numbers: $\sqrt{3}$

A.1.8.

$\sqrt{5}, -7, -\frac{7}{3}, 0, 3.14, \frac{5}{4}, -3, 12, 5$

- (a) Natural numbers: 12, 5
- (b) Whole numbers: 0, 12, 5
- (c) Integers: $-7, 0, -3, 12, 5$
- (d) Rational numbers: $-7, -\frac{7}{3}, 0, 3.14, \frac{5}{4}, -3, 12, 5$
- (e) Irrational numbers: $\sqrt{5}$

A.1.9.

2.01, $0.\overline{6}$, -13 , $0.010110111\dots$, 1 , -6

(a) Natural numbers: 1

(b) Whole numbers: 1

(c) Integers: -13 , 1 , -6

(d) Rational numbers: 2.01 , $0.\overline{6}$, -13 , 1 , -6

(e) Irrational numbers: $0.010110111\dots$

A.1.10.

25 , -17 , $-\frac{12}{5}$, $\sqrt{9}$, 3.12 , $\frac{1}{2}\pi$, 18 , -11.1 , 13

(a) Natural numbers: 25 , $\sqrt{9} = 3$, 18 , 13

(b) Whole numbers: 25 , $\sqrt{9} = 3$, 18 , 13


(c) Integers: 25 , -17 , $\sqrt{9} = 3$, 18 , 13

(d) Rational numbers: 25 , -17 , $-\frac{12}{5}$, $\sqrt{9} = 3$, 3.12 , 18 , -11.1 , 13

(e) Irrational numbers: $\frac{1}{2}\pi$


A.1.11.

$$\frac{5}{6} > \frac{2}{3}$$




A.1.12.

$$-4 > -8$$




A.1.13.

$$-5.2 > -8.5$$



A.1.14.

$$-\frac{4}{3} > -4.75$$



A.1.15.

The inequality $x \leq 5$ denotes the set of all real numbers less than or equal to 5.

A.1.16.

The inequality $x < 0$ denotes the set of all real numbers less than zero.

A.1.17.


The inequality $-2 < x < 2$ denotes the set of all real numbers greater than -2 and less than 2 .

A.1.18.

The inequality $0 < x \leq 6$ denotes the set of all positive real numbers less than or equal to 6 .


A.1.19.

The interval $[4, \infty)$ denotes the set of all real numbers greater than or equal to 4 ; $x \geq 4$.




A.1.20.

The interval $(-\infty, 2)$ denotes the set of all real numbers less than 2 ; $x < 2$.




A.1.21.

The interval $[-5, 2)$ denotes the set of all real numbers greater than or equal to -5 and less than 2 ; $-5 \leq x < 2$.




A.1.22.

The interval $(-1, 2]$ denotes the set of all real numbers greater than -1 and less than or equal to 2 ; $-1 < x \leq 2$.




A.1.23.

$$(-\infty, 0]; y \leq 0$$




A.1.24.

$$(-\infty, 25]; y \leq 25$$




A.1.25.

$$[10, 22]; 10 \leq t \leq 22$$



A.1.26.

$$[-3, 5); -3 \leq k < 5$$



A.1.27.

$$|-10| = -(-10) = 10$$

A.1.28.

$$|0| = 0$$

A.1.29.

$$|-1| - |-2| = 1 - 2 = -1$$

A.1.30.

$$-3 - |-3| = -3 - (3) = -6$$

A.1.31.

$$5 |-5| = 5(5) = 25$$

A.1.32.

$$-4 |-4| = -4(4) = -16$$

A.1.33.

If $x < -2$, then $x + 2$ is negative.

$$\text{So, } \frac{|x+2|}{x+2} = \frac{-(x+2)}{x+2} = -1.$$

A.1.34.

If $x > 1$, then $x - 1$ is positive.

$$\text{So, } \frac{|x-1|}{x-1} = \frac{x-1}{x-1} = 1.$$

A.1.35.

$$|-4| = |4| \text{ because } |-4| = 4 \text{ and } |4| = 4.$$

A.1.36.

$$-5 = -|5| \text{ because } -5 = -5.$$

A.1.37.

$$-|-6| < |-6| \text{ because } |-6| = 6 \text{ and } -|-6| = -(6) = -6.$$

A.1.38.

$$-|-2| = -|2| \text{ because } -2 = -2.$$

A.1.39.

$$d(126, 75) = |75 - 126| = 51$$

A.1.40.

$$d(-20, 30) = |30 - (-20)| = |50| = 50$$

A.1.41.

$$d\left(-\frac{5}{2}, 0\right) = \left|0 - \left(-\frac{5}{2}\right)\right| = \frac{5}{2}$$

A.1.42.

$$d\left(-\frac{1}{4}, -\frac{11}{4}\right) = \left|-\frac{11}{4} - \left(-\frac{1}{4}\right)\right| = \left|-\frac{5}{2}\right| = \frac{5}{2}$$

A.1.43.

$7x + 4$
Terms: $7x$, 4
Coefficient: 7

A.1.44.

$6x^3 - 5x$
Terms: $6x^3$, $-5x$
Coefficients: 6 , -5

A.1.45.

$4x^3 + 0.5x - 5$
Terms: $4x^3$, $0.5x$, -5
Coefficients: 4 , 0.5

A.1.46.

$3\sqrt{3}x^2 + 1$
Terms: $3\sqrt{3}x^2$, 1
Coefficient: $3\sqrt{3}$

A.1.47.

Receipts: $R = \$3268.0$ billion
Expenditures: $E = \$3852.6$ billion
 $|R - E| = |3268.0 - 3852.6| = |-584.6| = \584.6 billion

A.1.48.

Receipts: $R = \$3329.9$ billion
Expenditures: $E = \$4109.0$ billion
 $|R - E| = |3329.9 - 4109.0| = |-779.1| = \779.1 billion

A.1.49.

Receipts: $R = \$3421.2$ billion
Expenditures: $E = \$6553.6$ billion
 $|R - E| = |3421.2 - 6553.6| = |-3132.4| = \3132.4 billion

A.1.50.

Receipts: $R = \$4897.3$ billion

Expenditures: $E = \$6273.3$ billion

$$|R - E| = |4897.3 - 6273.3| = |-1376.0| = \$1376.0 \text{ billion}$$

A.1.51.

$$x^2 - 3x + 2$$

$$(a) (0)^2 - 3(0) + 2 = 2$$

$$(b) (-1)^2 - 3(-1) + 2 = 1 + 3 + 2 = 6$$

A.1.52.

$$\frac{x-2}{x+2}$$

$$(a) \frac{2-2}{2+2} = \frac{0}{4} = 0$$

$$(b) \frac{-2-2}{-2+2} = \frac{-4}{0}$$

Division by zero is undefined.

A.1.53.

$$\frac{2x}{3} - \frac{x}{4} = \frac{8x}{12} - \frac{3x}{12} = \frac{5x}{12}$$

A.1.54.

$$\frac{3x}{4} + \frac{x}{5} = \frac{15x}{20} + \frac{4x}{20} = \frac{19x}{20}$$

A.1.55.

$$\frac{3x}{10} \cdot \frac{5}{6} = \frac{x}{2} \cdot \frac{1}{2} = \frac{x}{4}$$

A.1.56.

$$\frac{2x}{3} \div \frac{6}{7} = \frac{2x}{3} \cdot \frac{7}{6} = \frac{x}{3} \cdot \frac{7}{3} = \frac{7x}{9}$$

A.1.57.

False. Because zero is nonnegative but not positive, not every nonnegative number is positive.

A.1.58.

The product of two negative numbers is positive.

A.1.59.

The distributive law was used incorrectly. The 5 was not distributed to the 3.

A.1.60.

- (a) Because the price can only be a positive rational number with at most two decimal places, the description matches graph (ii).

A range of prices can only include zero and positive numbers with at most two decimal places. So, a range of prices can be represented by whole numbers and some noninteger positive fractions.

- (b) Because the distance is a positive real number, the description matches graph (i).

A range of lengths can only include positive numbers. So, a range of lengths can be represented by positive real numbers.

A.1.61.

$5/n$ increases or decreases without bound. As n approaches 0, $5/n$ increases when n is positive and decreases when n is negative.

A.1.62.

$5/n$ approaches 0 as n increases without bound. As n increases, $5/n$ decreases but stays positive.

A.1.63.

$$6 = 2 \cdot 3, 8 = 2 \cdot 2 \cdot 2$$

- (a) Greatest common factor: 2

- (b) Least common multiple: $2 \cdot 2 \cdot 2 \cdot 3 = 24$

A.1.64.

$$10 = 2 \cdot 5, 25 = 5 \cdot 5$$

- (a) Greatest common factor: 5

- (b) Least common multiple: $2 \cdot 5 \cdot 5 = 50$

A.1.65.

$$27 = 3 \cdot 3 \cdot 3, 36 = 2 \cdot 2 \cdot 3 \cdot 3, 54 = 2 \cdot 3 \cdot 3 \cdot 3$$

- (a) Greatest common factor: $3 \cdot 3 = 9$

- (b) Least common multiple: $2 \cdot 2 \cdot 3 \cdot 3 \cdot 3 = 108$

A.1.66.

$$49 = 7 \cdot 7, 98 = 2 \cdot 7 \cdot 7, 112 = 2 \cdot 2 \cdot 2 \cdot 7$$

- (a) Greatest common factor: 7

- (b) Least common multiple: $2 \cdot 2 \cdot 2 \cdot 2 \cdot 7 \cdot 7 = 784$

A.1.67.

$$-(3 \cdot 3) = -9$$

A.1.68.

$$(-3)(-3) = 3 \cdot 3 = 9$$

A.1.69.

$$\frac{(-5)(-5)}{-5} = -5$$

A.1.70.

$$\frac{-(5 \cdot 5)}{(-5)(-5)(-5)} = \frac{5 \cdot 5}{5 \cdot 5 \cdot 5} = \frac{1}{5}$$

A.1.71.

$$48 = 2 \cdot 2 \cdot 2 \cdot 2 \cdot 3 = 2^4 \cdot 3$$

A.1.72.

$$250 = 2 \cdot 5 \cdot 5 \cdot 5 = 2 \cdot 5^3$$

A.1.73.

$$792 = 8 \cdot 9 \cdot 11 = 2 \cdot 2 \cdot 2 \cdot 3 \cdot 3 \cdot 11 = 2^3 \cdot 3^2 \cdot 11$$

A.1.74.

$$4802 = 2 \cdot 7 \cdot 7 \cdot 7 \cdot 7 = 2 \cdot 7^4$$

A.1.75.

$$3.785(10,000) = 37,850$$

A.1.76.

$$1.42(1,000,000) = 1,420,000$$

A.1.77.

$$\frac{6.09}{1000} = 0.00609$$

A.1.78.

$$\frac{8.603}{100,000} = 0.00008603$$

APPENDIX A.2 EXPONENTS AND RADICALS

A.2.1.

exponent; base

A.2.2.

index; radicand

A.2.3.

An expression involving radicals is in simplest form when the following conditions are satisfied:

1. All possible factors have been removed from the radical.
2. All fractions have radical-free denominators.
3. The index of the radical is reduced.

A.2.4.

64 is both a perfect square ($8^2 = 64$) and a perfect cube ($4^3 = 64$).

A.2.5.

$$5 \cdot 5^3 = 5^4 = 625$$

A.2.6.

$$\begin{aligned} (2^3 \cdot 3^2)^2 &= 2^{3 \cdot 2} \cdot 3^{2 \cdot 2} \\ &= 2^6 \cdot 3^4 = 64 \cdot 81 = 5184 \end{aligned}$$

A.2.7.

$$\frac{5^2}{5^4} = 5^{-2} = \frac{1}{5^2} = \frac{1}{25}$$

A.2.8.

$$\begin{aligned} \left(-\frac{3}{5}\right)^3 \left(\frac{5}{3}\right)^2 &= (-1)^3 \frac{3^3}{5^3} \cdot \frac{5^2}{3^2} = -1 \cdot 3^{3-2} \cdot 5^{2-3} \\ &= -3 \cdot 5^{-1} = -\frac{3}{5} \end{aligned}$$

A.2.9.

$$-3^2 = -9$$

A.2.10.

$$(-4)^{-3} = \frac{1}{(-4)^3} = -\frac{1}{64}$$

A.2.11.

$$\frac{4 \cdot 3^{-2}}{2^{-2} \cdot 3^{-1}} = 4 \cdot 2^2 \cdot 3^{-2-(-1)} = 4 \cdot 4 \cdot 3^{-1} = \frac{16}{3}$$

A.2.12.

$$\frac{3}{3^{-4}} = 3 \cdot 3^4 = 3^5 = 243$$

A.2.13.

When $x = 2$,

$$-3x^3 = -3(2)^3 = -24.$$

A.2.14.

When $x = 4$,

$$7x^{-2} = 7(4)^{-2} = \frac{7}{4^2} = \frac{7}{16}.$$

A.2.15.

When $x = 0.1$,

$$6x^2 = 6(0.1)^2 = 6(0.01) = 0.06.$$

A.2.16.

When $x = -\frac{1}{3}$,

$$12(-x)^3 = 12\left(\frac{1}{3}\right)^3 = 12\left(\frac{1}{27}\right) = \frac{4}{9}.$$

A.2.17.

$$6y^2(2y^0)^2 = 6y^2(2 \cdot 1)^2 = 6y^2(4) = 24y^2$$

A.2.18.

$$\begin{aligned} (-z)^3(3z^4) &= (-1)^3(z^3)3z^4 \\ &= -1 \cdot 3 \cdot z^{3+4} = -3z^7 \end{aligned}$$

A.2.19.

$$\frac{7x^2}{x^3} = 7x^{2-3} = 7x^{-1} = \frac{7}{x}$$